


Elastic collision of two bright solitons, $|\psi(x,t)|^2$, by split-step Fourier.

Numerical Methods for the Nonlinear Schrödinger Equation

A Method of Lines Approach for the NLSE and GPE with Python

Þráinn Eiríksson



ODIN PRESS



Numerical Methods for the Nonlinear Schrödinger Equation

© 2026 **Þráinn Eiríksson**. All Rights Reserved.

All rights reserved. No part of this book — including text, equations, figures, and the Python source code listings contained herein — may be reproduced, stored in a retrieval system, or transmitted in any form or by any means (electronic, mechanical, photocopying, recording, or otherwise) without the prior written permission of the author, except in the case of brief quotations embodied in critical reviews and certain other noncommercial uses permitted by copyright law. For permission requests, contact the author directly.

The Python source code presented in this book is original work by the author. It may be used for personal study and non-commercial research purposes, provided that the source is acknowledged. Any other use requires written permission from the author.

Every effort has been made to ensure the accuracy of the mathematical content. The author accepts no liability for errors or omissions.

Published by Odin Press.
`https://odinpress.org`
This book was typeset in LaTeX in Latin Modern Roman by Odin Press from a manuscript submitted by the author.
Paperback version, trim size 6 × 9 inches.
ISBN: forthcoming (Bowker allocation pending)

9 8 7 6 5 4 3 2 1 0

Preface

The nonlinear Schrödinger equation is one of the most successful models in mathematical physics. It governs the envelope of a weakly nonlinear dispersive wave in one spatial dimension, appearing in contexts as different as optical fibre communication, Bose–Einstein condensates, deep-water wave propagation, and plasma physics. Its one-dimensional defocusing cousin, the Gross–Pitaevskii equation, describes superfluid dynamics in a dilute atomic gas. What makes both equations remarkable is not their breadth of application but their depth of mathematical structure: they are exactly solvable by the inverse scattering transform, and their solutions — solitons, breathers, rogue waves, cnoidal waves — can be written down in closed form.

This book is about computing those solutions numerically, and computing them well. The approach throughout is the Method of Lines: the spatial variable is discretised by a pseudospectral operator (DST-I spline collocation for problems with Dirichlet boundary conditions, Fourier pseudospectral for problems with periodic ones), leaving an autonomous system of ordinary differential equations that is integrated in time by an explicit adaptive Runge–Kutta solver from SciPy. Every chapter applies this pipeline to a different exact solution, verifies the numerical output against the closed form to pointwise accuracy 10^{-10} or better, and monitors the conserved quantities — L^2 norm, momentum, and Hamiltonian energy — as independent diagnostics. The result, ac-

cumulated over fourteen chapters, is a complete computational atlas of the NLS/GPE solution zoo.

The book is organised in a single progression. Chapters 1 through 3 treat the elementary localised solutions of the focusing and defocusing NLS: the bright soliton, the black soliton, and the grey soliton family. Chapters 4 and 5 introduce the Darboux transformation and use it to construct multi-soliton exact solutions: the elastic two-soliton collision and the two-soliton bound state (soliton molecule). Chapters 6 and 7 leave the soliton family and treat the spatially periodic cnoidal waves via Jacobi elliptic functions, first with Dirichlet and then with periodic boundary conditions. Chapter 8 studies modulational instability — the Benjamin–Feir mechanism by which a plane wave amplifies sidebands exponentially — and introduces a different verification strategy based on linear growth rates rather than a pointwise exact solution. Chapters 9 through 11 treat the rogue-wave hierarchy sitting on a plane-wave background: the Akhmediev breather (periodic in space), the Kuznetsov–Ma breather (periodic in time), and the Peregrine soliton (localised in both variables), the latter being the degenerate limit of the former two. Chapter 12 extends the pipeline to the Manakov vector NLS, showing how the two-component system inherits elastic collisions and exact solutions from the scalar theory. Chapter 13 constructs the three-soliton collision via a three-step Darboux transform and measures the pairwise phase shifts analytically. Chapter 14 closes Part I with the second-order rational rogue wave, the $N = 2$ member of the Peregrine hierarchy, whose peak amplitude is five times the background. Chapter 15 is an epilogue: it explains the mathematical reasons the pipeline works, identifies its honest limits, and sketches ten natural extensions that go beyond the exact-solution regime.

The computational tools are deliberately minimal. No specialised NLS library is required. The only dependencies are NumPy, SciPy, and Matplotlib, all part of any standard scientific Python installation.

Every solver is a self-contained script of fewer than one hundred lines; every figure is generated by a companion script that can be run independently. The goal is transparency: a reader who finishes the book should be able to reproduce any result from scratch and extend any solver to a new problem without needing to understand a large pre-existing codebase.

The intended audience is graduate students and researchers in applied mathematics, theoretical physics, or photonics who are comfortable with ordinary differential equations and Fourier analysis, and who want a practical introduction to spectral numerical methods for dispersive nonlinear PDEs. The book assumes no prior knowledge of soliton theory; the necessary exact solutions are derived or cited from the primary literature as they are needed. It equally assumes no prior knowledge of the Method of Lines; the spatial discretisation and its accuracy theory are developed from first principles in Chapter 1 and carried forward unchanged.

A note on verification philosophy: every chapter in this book was written after the solver was verified, not before. The exact solution comes first; the numerical output is compared against it before a single sentence of text is written. This discipline — exact solution, verified solver, then explanation — is the book's only methodology. It is also, the author believes, the right way to write about numerical methods.

Þráinn Eiríksson
2026

Contents

Preface **vii**

1 The Focusing NLS — Bright Soliton **1**
1.1 Introduction 1
1.2 The Nonlinear Schrödinger Equation 2
1.2.1 Dispersion Relation 3
1.2.2 The Real–Imaginary Splitting 3
1.3 The Exact One-Soliton Solution 3
1.3.1 Special Cases 5
1.4 The Hamiltonian Structure and Conservation Laws . . 5
1.5 Spline Collocation Discretisation 7
1.5.1 The MOL Semi-Discretisation 8
1.5.2 Spectral Convergence 9
1.6 Discrete Conservation Laws 10
1.7 Why RK45 and Not Radau 11
1.8 Implementation 12
1.9 Numerical Results 13

2 The Defocusing NLS — Dark Soliton **17**
2.1 Introduction 17
2.2 The Defocusing NLS 18
2.2.1 Conservation Laws 19

2.3 The Dark Soliton 20
2.3.1 Exact Formula 20
2.3.2 Density Profile 22
2.3.3 Phase Structure 22
2.4 Smoothness 23
2.5 Boundary Conditions — Linear-Lift Correction 24
2.5.1 Non-Zero Boundary Values 24
2.5.2 The Linear-Lift Lemma 24
2.5.3 Accuracy of the Correction 25
2.6 Implementation 25
2.7 Spectral Convergence 28
2.8 Numerical Results 29
2.8.1 Density and Phase 29
2.8.2 Pointwise Error 30
2.8.3 Norm Conservation 31
2.9 Summary 31

3 The Grey Soliton — Defocusing NLS with Velocity 33
3.1 Introduction 33
3.2 The Defocusing NLS and Background States 35
3.2.1 Conservation Laws 35
3.2.2 Plane-Wave Background 36
3.3 The Grey Soliton 36
3.3.1 Exact Formula 36
3.3.2 Density Profile 38
3.3.3 Phase Structure 38
3.4 Velocity, Darkness, and the Soliton Family 39
3.4.1 The One-Parameter Family 39
3.4.2 Conservation of the Notch Intensity 40
3.5 Smoothness and Spectral Approximability 40
3.6 Boundary Conditions and Linear-Lift Correction . . . 41
3.7 Implementation 42

3.8 Spectral Convergence 45
3.9 Numerical Results 46
3.9.1 Density and Phase 46
3.9.2 Pointwise Error 47
3.9.3 Norm Conservation 48
3.9.4 Comparison with the Dark Soliton 48
3.10 Summary 49

4 Elastic Two-Soliton Collision — Focusing NLS 51
4.1 Introduction 51
4.2 The Focusing NLS and the Two-Soliton Problem . . . 52
4.2.1 Real–Imaginary Splitting 53
4.2.2 The Two-Soliton Initial Condition 53
4.3 The Darboux Exact Two-Soliton Solution 54
4.3.1 The Zakharov–Shabat Scattering Problem . . . 54
4.3.2 Vacuum Eigenfunctions 54
4.3.3 Two Darboux Steps from the Vacuum 55
4.4 The Elastic Collision and Phase Shift 56
4.4.1 Asymptotic Analysis 56
4.4.2 The Phase Shift Formula 57
4.5 Conservation Laws 58
4.6 Spline Collocation Discretisation 60
4.6.1 DST-I Pseudospectral Operator 60
4.6.2 The MOL System 60
4.6.3 Choice of Integrator 61
4.7 Numerical Verification 61
4.7.1 Verification Table 61
4.7.2 The Error Spike at the Collision 62
4.8 Implementation 62
4.9 Numerical Results 65
4.10 Discussion 69

5 Soliton Molecules — Two-Soliton Bound State **71**
5.1 Introduction . . . 71
5.2 The Focusing NLS and the Bound-State Problem . . . 72
5.2.1 Real–Imaginary Splitting . . . 73
5.2.2 Parameters of the Bound State . . . 73
5.3 The ZS Binding Condition . . . 74
5.4 The Exact Two-Soliton Bound State . . . 75
5.5 Breathing Period and Peak Amplitude . . . 77
5.6 Conservation Laws . . . 78
5.7 Spline Collocation Discretisation . . . 79
5.7.1 Integrator . . . 80
5.7.2 Hamiltonian Discretisation . . . 80
5.8 Numerical Verification . . . 81
5.9 Implementation . . . 82
5.10 Numerical Results . . . 86
5.11 Discussion . . . 89

6 Cnoidal Wave — Jacobi Elliptic Solution of Focusing NLS **91**
6.1 Introduction . . . 91
6.2 The Focusing NLS and the Cnoidal Wave Problem . . . 93
6.2.1 Real–Imaginary Splitting . . . 94
6.2.2 Problem Parameters . . . 94
6.3 Jacobi Elliptic Functions — Definitions and Key Identities . . . 95
6.4 The Exact Cnoidal Wave Solution . . . 96
6.4.1 Properties of the Exact Solution . . . 97
6.5 Domain Choice and Exact Boundary Conditions . . . 97
6.5.1 Why DST-I and Not Fourier Spectral? . . . 98
6.6 Limiting Cases . . . 99
6.6.1 Soliton Limit . . . 99
6.6.2 Linear Wave Limit . . . 100

6.7 Conservation Laws 100
6.7.1 Exact Norm Formula 100
6.7.2 Hamiltonian 101
6.7.3 Discrete Norm Conservation 102
6.8 Spatial Discretisation 102
6.8.1 Resolution Analysis 102
6.8.2 N=64 is Optimal, Not Just Sufficient 103
6.9 Numerical Verification 103
6.10 Implementation 104
6.11 Numerical Results 110
6.11.1 Spacetime Density — Figure 6.1 110
6.11.2 Spatial Profile and Pointwise Error — Figure 6.2 111
6.11.3 Conservation Diagnostics — Figure 6.3 112
6.11.4 Parameter Study — Figure 6.4 113
6.12 Discussion 114
6.12.1 The Cnoidal Wave in the IST Hierarchy 114
6.12.2 Relationship to Modulational Instability 114
6.12.3 Preview of Chapter 7 115

7 Cnoidal Wave — Jacobi dn Solution of Focusing NLS 117
7.1 Introduction 117
7.2 The dn Wave Problem 119
7.2.1 Why periodic, not Dirichlet? 119
7.2.2 Contrast with Chapter 6 120
7.3 The Jacobi dn Function 120
7.3.1 Definition and basic properties 120
7.3.2 The second-derivative identity 121
7.4 The Exact dn Wave Solution 122
7.5 Limiting Cases 123
7.5.1 Rate of convergence to the soliton limit 124
7.6 Conservation Laws 125
7.6.1 Exact norm 125

7.6.2 Discrete norm conservation 126
7.6.3 Hamiltonian 127
7.7 Orbital Stability 127
7.8 Fourier Pseudospectral Discretisation 129
7.8.1 Grid and operator 129
7.8.2 Comparison with DST-I from Chapter 6 130
7.8.3 Resolution analysis 130
7.9 Numerical Verification 130
7.10 Orbital Stability Experiment 132
7.11 Parameter Study 133
7.12 Implementation 133
7.13 Numerical Results 139
7.14 Discussion 140

8 Modulational Instability — The Benjamin–Feir Instability 145
8.1 Introduction 145
8.2 Plane-Wave Solution 146
8.2.1 The PDE and Exact Plane-Wave Family 146
8.2.2 Physical Interpretation 147
8.3 Linear Stability Analysis 148
8.3.1 Linearised Equations 148
8.3.2 Derivation of the Dispersion Relation 148
8.3.3 The Benjamin–Feir Criterion 149
8.4 Growth Rate and Instability Spectrum 150
8.4.1 The Growth Rate Formula 150
8.4.2 The Instability Window 151
8.4.3 Integer Mode Selection and Domain Choice . . 151
8.5 Conservation Laws 152
8.5.1 The L^2 Norm 152
8.5.2 The Hamiltonian 152
8.5.3 Conservation Under the NLS Flow 152

8.6 Fourier Pseudospectral Method of Lines 153
8.6.1 The Fourier D^2 Operator 153
8.6.2 The MOL Right-Hand Side and Time Integration 154
8.6.3 Optimality of the Fourier Method for This Problem 154
8.7 Numerical Verification 155
8.7.1 Parameters and Mode Extraction 155
8.7.2 Verification Results 155
8.8 Spacetime Evolution 156
8.9 Growth Rate Verification 157
8.10 Growth Rate Spectrum 158
8.11 Conservation Diagnostics 159
8.12 Effect of Perturbation Amplitude 160
8.13 Connection to Nonlinear Theory 161
8.14 Summary 162

9 Akhmediev Breather — Spatially Periodic Soliton on a Plane-Wave Background 163
9.1 Introduction 163
9.2 The Soliton-on-Finite-Background Family 165
9.2.1 A Unified Parametrisation 165
9.2.2 The Kuznetsov–Ma and Peregrine Formulas . . 165
9.2.3 Why the Akhmediev Branch Is Optimal for Fourier MOL 166
9.3 Exact Formula and Properties 166
9.3.1 Parameters and Notation 166
9.3.2 Numerator Decomposition 167
9.3.3 The Exact Solution Theorem 168
9.3.4 Denominator Positivity 169
9.3.5 Return to Background 170
9.3.6 Peak Amplitude 170
9.4 Connection to Benjamin–Feir Theory 171

9.4.1 Growth Rate Identity 171
9.4.2 The Peregrine Limit 172
9.5 Fourier Pseudospectral Discretisation 172
9.6 Implementation 173
9.7 Numerical Verification 180
9.8 Numerical Results 181
9.8.1 Spacetime Density — Figure 9.1 181
9.8.2 Density Profiles at Fixed Times — Figure 9.2 . 182
9.8.3 Parameter Study — Figure 9.3 183
9.8.4 Growth Rate Verification — Figure 9.4 184
9.8.5 Conservation Diagnostics — Figure 9.5 185
9.9 Discussion 186

10 Kuznetsov–Ma Breather — Temporally Periodic Soliton on a Plane-Wave Background 189
10.1 Introduction 189
10.2 The Kuznetsov–Ma Solution 191
10.2.1 Formula and Parameters 191
10.2.2 Numerator Decomposition 192
10.2.3 Contrast with the Akhmediev Breather 193
10.2.4 Reference Values 193
10.3 Exact Solution Theorem 193
10.4 Background Return 196
10.5 Peak Amplitude 197
10.6 Peregrine Limit 198
10.7 Excess Norm 200
10.8 Fourier Pseudospectral Discretisation 202
10.8.1 Why Fourier for a Plane-Wave Background . . 202
10.8.2 Haragus–Pelinovsky Normalisation vs Book Normalisation 203
10.8.3 Grid and the Aliasing Bottleneck 203
10.8.4 Integrator 205

10.9 Python Implementation 206
10.10 Numerical Verification and Results 213
10.10.1 Reference Run 213
10.10.2 Six-Point Pointwise Error 213
10.10.3 Conservation 213
10.10.4 Parameter Study 214
10.10.5 Figures 214
10.11 Discussion and Outlook 218

11 Peregrine Soliton — Doubly-Localised Rational Rogue Wave 221
11.1 Introduction 221
11.2 The Peregrine Solution 224
11.2.1 Formula and Parameters 224
11.2.2 Key Exact Properties 225
11.2.3 Contrast with Akhmediev and Kuznetsov–Ma . 225
11.3 Derivation and Verification of the Exact Solution . . . 226
11.3.1 Rational Ansatz 226
11.3.2 Direct Substitution 227
11.4 Global Amplitude Bound 229
11.5 Algebraic Decay and Boundary Residual 231
11.6 Conservation Laws 233
11.7 Modulational Instability and the Double-Precision Floor 234
11.7.1 Linear Growth Rate on the Plane-Wave Background 235
11.7.2 Quantitative Floor 236
11.7.3 Why the Floor Is Not Method-Dependent . . . 237
11.8 Failed Methods 238
11.8.1 DST-I (Dirichlet Sine Transform) with Co-Rotating Lift 238
11.8.2 Finite Differences (High-Order Central Stencils) 239
11.8.3 Chebyshev Pseudospectral 239

11.8.4 Radau (Implicit Integrator) 239
11.8.5 Yoshida-4 Symplectic 240
11.8.6 Lawson–RK4 Exponential Integrator 240
11.8.7 Method Selection Summary 241
11.9 The Successful Method 242
11.9.1 Fourier $\mathcal{D}_2$ 242
11.9.2 DOP853 242
11.9.3 Split at $t = 0$ 243
11.9.4 Discrete Conservation 243
11.10 Python Implementation 244
11.10.1 Solver 244
11.10.2 Figures 247
11.11 Numerical Verification 252
11.11.1 Reference Run 252
11.11.2 Pointwise Error at Six Test Times 252
11.11.3 Conservation Diagnostics 253
11.11.4 Peak Amplitude 253
11.12 Figures 253
11.13 Discussion and Outlook 258

12 Manakov System — Vector NLS and the Polarisation-Rotating Collision 261
12.1 Introduction 261
12.2 The Manakov System and Its Integrability 264
12.2.1 Governing System and Hamiltonian Structure . 264
12.2.2 Conservation Laws 264
12.2.3 Lax Pair and Integrability 266
12.2.4 Well-Posedness 266
12.3 The One-Soliton Solution 267
12.4 The Two-Soliton Bilinear Solution 268
12.4.1 Statement of the Formula 268
12.4.2 Initial-Centre Calibration 270

12.4.3 Orthogonal-Polarisation Simplification 271
12.4.4 Numerical Values for the Chapter's Configuration 271
12.5 Polarisation Transformation Across the Collision . . . 271
12.5.1 Asymptotic Separation 271
12.6 Spatial Discretisation for the Vector Problem 273
12.6.1 Four-Real-Component State Vector 273
12.6.2 Right-Hand Side 273
12.6.3 Semi-Discrete Conservation 274
12.7 Python Implementation 275
12.8 Numerical Verification 280
12.8.1 Parameters 280
12.8.2 Pointwise Error 280
12.8.3 Conservation Diagnostics 281
12.8.4 Diagnostic Figures 282
12.9 Discussion and Outlook 286

13 Elastic Three-Soliton Collision **287**
13.1 Introduction 287
13.2 The Focusing Nonlinear Schrödinger Equation 288
13.3 Soliton Parameters and the Zakharov–Shabat Eigenvalue 290
13.4 Seed Eigenfunctions 291
13.5 The Darboux Transformation and the Three-Soliton Solution 291
13.5.1 Single Darboux Step 291
13.5.2 Three-Step Construction 292
13.5.3 Peak Amplitude at the Triple Collision 293
13.6 Conservation Laws and Their Numerical Verification . 294
13.7 Elastic Collision and Phase Shifts 295
13.7.1 Shape Preservation 295
13.7.2 Phase Shift Formula 295
13.8 Spatial Discretisation: DST-I Pseudospectral Method 296
13.8.1 Grid Resolution Analysis 298

13.9 Time Integration 298
13.10 Solver Verification 299
13.11 Numerical Results 299
13.12 Implementation 303
13.13 Discussion 309

14 Second-Order Rogue Wave **311**
14.1 Introduction 311
14.2 The Focusing NLS and the Plane-Wave Background . 313
14.3 Rational Rogue-Wave Hierarchy — Darboux Construction 314
14.4 The Second-Order Rogue Wave: Exact Formula 315
14.5 Conservation Laws 317
14.6 Spatial Structure: Satellite Peaks and Triangular Pattern 318
14.7 Modulational Instability of the Background 320
14.8 Fourier Pseudospectral Discretisation 321
14.8.1 Split-at-$t = 0$ Strategy 322
14.8.2 Time Integrator 322
14.9 Solver Verification 323
14.10 Numerical Results 324
14.11 Comparison with the Peregrine Soliton ($N = 1$) 327
14.12 Implementation 328
14.13 Discussion 331

15 Epilogue — The Pipeline, Its Foundations, and the Road Ahead **333**
15.1 Introduction 333
15.2 The Solution Zoo — A Synthesis of Chapters 1–14 . . 334
15.3 The Mathematical Pillars 336
15.3.1 Complete Integrability and the Inverse Scattering Transform 336

15.3.2 The Darboux Transformation — An Exact-Solution Factory 338
15.3.3 Discrete Conservation Laws and the Symmetry of the Spatial Operator 339
15.3.4 Analyticity and Spectral Accuracy — The Single Theorem Behind All Fourteen Chapters . . 341
15.4 The Road Ahead — Ten Extensions 342
15.4.1 Delta-Function Potential — Soliton Scattering 343
15.4.2 Double-Well Trap — Josephson Oscillations and Self-Trapping 344
15.4.3 Optical Lattice — Bloch States and Gap Solitons 345
15.4.4 Saturable Nonlinearity — Photorefractive Solitons 346
15.4.5 Cubic-Quintic NLS — Competing Nonlinearities and Collapse Arrest 347
15.4.6 Nonlocal NLS — Liquid Crystal Response . . . 347
15.4.7 Two-Component BEC — Phase Separation . . 348
15.4.8 Spin-Orbit Coupled BEC — Synthetic Gauge Fields 349
15.4.9 Anderson Localisation — Random Potentials . 350
15.4.10 Two-Dimensional GPE — Vortices, Collapse, and the Townes Threshold 351
15.5 Limits of the Pipeline 352
15.6 Closing Remarks 353

References **357**

Chapter 1

The Focusing NLS — Bright Soliton

1.1 Introduction

The nonlinear Schrödinger equation (NLS) is one of the most pervasive equations in mathematical physics. It governs the propagation of optical pulses in nonlinear fibres, the dynamics of Bose–Einstein condensates, and the modulation of water waves in deep water. The equation was solved exactly by Zakharov and Shabat [65] using the inverse scattering method, establishing the NLS as a completely integrable Hamiltonian system with an infinite family of conservation laws.

This chapter introduces the full computational pipeline used throughout the book. We study the bright one-soliton — the simplest exact solution — and verify the numerical method against it at machine precision. The spatial operator is a not-a-knot cubic spline collocation second-derivative, following the framework developed by de Boor [14]. Time integration uses the explicit Runge–Kutta RK45 scheme, which is well-suited to the Hamiltonian structure of the NLS: the equation is not stiff, and explicit methods preserve the symplectic

phase-space geometry far better than implicit schemes for long-time integration. The method of lines (MOL) framework within which all this sits is due to Schiesser [54].

The pipeline established here — spline collocation, Re/Im splitting, RK45 integration, and conservation-law diagnostics — is reused unchanged in every subsequent chapter. Mastering it here removes all technical friction from the chapters that follow.

1.2 The Nonlinear Schrödinger Equation

Notation 1.1 (Complex wave function)**.** Throughout this chapter $\psi\colon [-L, L] \times [0, T] \to \mathbb{C}$ denotes the complex wave function. We write $u = \mathrm{Re}(\psi)$ and $v = \mathrm{Im}(\psi)$, so that $\psi = u + iv$. The modulus is $|\psi|^2 = u^2 + v^2$ and the L^2 norm is $\|\psi(t)\|^2 = \int_{-L}^{L} |\psi(x,t)|^2\, dx$.

Assumption 1.2 (Domain)**.** The spatial domain is $\Omega = [-L, L]$ with $L = 30$. The domain is chosen large enough that the soliton tails satisfy $|\psi(\pm L, t)| < 10^{-14}$ for all $t \in [0, T]$, so that homogeneous Dirichlet boundary conditions introduce no appreciable error.

Definition 1.3 (Focusing Cubic NLS)**.** The *focusing cubic nonlinear Schrödinger equation* is

$$i\,\partial_t \psi + \partial_{xx}\psi + 2|\psi|^2\psi = 0, \quad x \in \Omega,\ t \in [0, T], \tag{1.1}$$

subject to initial condition $\psi(x, 0) = \psi_0(x)$ and zero boundary conditions $\psi(\pm L, t) = 0$.

Remark 1.4. Equation (13.1) follows the normalisation of Sulem and Sulem [57]. The factor 2 yields the cleanest one-soliton form. The positive sign of the cubic term indicates *focusing*: nonlinearity concentrates the wave and bright solitons persist. The *defocusing* variant reverses this sign and admits dark solitons on a nonzero background.

1.2.1 Dispersion Relation

Before introducing the exact solution, we record the dispersion relation of the linearised problem. Setting $|\psi|^2 = 0$ in (13.1), plane-wave solutions $e^{i(kx-\omega t)}$ satisfy

$$\omega = k^2. \tag{1.2}$$

This is the free-particle dispersion relation of quantum mechanics. Nonlinearity modifies it: a wave packet of wavenumber k acquires the nonlinear frequency shift $-2|\psi|^2$, so the full phase velocity of a soliton of amplitude η and velocity ξ is $\xi/2-2\eta^2/\xi$, which is amplitude-dependent.

1.2.2 The Real–Imaginary Splitting

We separate (13.1) into its real and imaginary parts. Substituting $\psi = u + iv$ into (13.1) and equating real and imaginary parts gives the coupled real system

$$\partial_t u = -\partial_{xx} v - 2(u^2 + v^2)v, \tag{1.3}$$

$$\partial_t v = \quad \partial_{xx} u + 2(u^2 + v^2)u. \tag{1.4}$$

Equations (1.3)–(1.4) form a real ODE system after spatial discretisation, which avoids complex arithmetic in the time integrator. This splitting is computationally advantageous: real-valued arithmetic is faster and numerically cleaner. Every theorem and diagnostic in this chapter applies to the coupled system (1.3)–(1.4).

1.3 The Exact One-Soliton Solution

The existence of exact soliton solutions is the defining feature of integrable systems. The NLS was shown to be integrable — and its

Chapter 3

The Grey Soliton — Defocusing NLS with Velocity

3.1 Introduction

Chapter 2 introduced the *dark soliton* of the defocusing nonlinear Schrödinger equation: a travelling density notch whose minimum intensity reaches exactly zero. That construction rested on the special case $v = 0$, in which the soliton is stationary and the background phase rotates uniformly. The dark soliton is, however, the boundary case of a continuous one-parameter family of exact solutions. When the velocity v is non-zero, the density notch no longer touches zero; instead it floats at a grey level above the background, giving rise to the name *grey soliton*.

The grey soliton was identified by Zakharov and Shabat [65] as part of the full inverse scattering solution for the defocusing NLS: every bound state of the associated scattering problem corresponds to a single grey soliton parameterised by a complex eigenvalue $\lambda = i\kappa + \sigma$,

where κ controls the background amplitude and σ controls the velocity. A systematic classification appears in Ablowitz and Segur [2]. The physical significance of grey solitons extends to repulsive Bose–Einstein condensates, where Frantzeskakis [23] reviews extensive experiments in which phase-engineering imprints dark and grey solitons into condensates and observes their velocity-dependent contrast.

The central parameter distinguishing grey from dark is the *darkness parameter* $B = \sqrt{1 - C^2}$, where $C = v/(2A)$ is the normalised velocity. For $C = 0$ ($v = 0$): $B = 1$ and the density dips to zero — the black (maximally dark) soliton. As $|C| \to 1$ ($|v| \to 2A$): $B \to 0$ and the density notch vanishes entirely; the grey soliton dissolves into a linear dispersive wave. Between these limits, the grey soliton traverses the domain at speed v while maintaining a non-zero intensity minimum A^2C^2 and a phase jump strictly less than π.

For this chapter we choose $A = 1$, $v = 1.2$, $x_0 = -6$, giving $C = 0.6$, $B = 0.8$, and a notch minimum $|\psi|^2_{\min} = 0.36$: the grey soliton retains 36
intensity at its centre. This is well within the *genuinely grey* regime, clearly distinguishable from the nearly-dark soliton of Chapter 2 ($v = 0.5$, minimum 6.25%). The soliton starts at $x_0 = -6$ and travels to $x \approx 6$ over $T = 10$ time units, remaining well inside the domain $(-30, 30)$ throughout.

The numerical method is identical to Chapter 2: DST-I spline collocation for ∂_{xx} with linear-lift correction for the non-zero boundary values, and RK45 with rtol $= 10^{-12}$, atol $= 10^{-14}$ from Dormand and Prince [19]. The grey soliton is C^∞, so the DST-I operator retains spectral accuracy. The maximum pointwise error at $T = 10$ with $N = 512$ is 2.52×10^{-13} — more than three orders of magnitude below the 10^{-10} target.

3.2 The Defocusing NLS and Background States

The governing equation is the defocusing cubic NLS, repeated here for self-containment:

$$i\partial_t \psi + \partial_{xx}\psi - 2|\psi|^2\psi = 0, \quad x \in (-L, L),\ t > 0. \tag{3.1}$$

Notation 3.1 (Chapter conventions)**.** All definitions and equation labels in this chapter are prefixed 3 to avoid confusion with Chapter 2. Equation (3.1) is identical to (2.1); results from Chapter 2 are cited directly where applicable.

3.2.1 Conservation Laws

Theorem 3.2 (Conservation of norm and energy)**.** *If ψ solves (3.1) with Dirichlet or periodic boundary conditions, then the L^2 norm*

$$\mathcal{N}[\psi] = \int_{-L}^{L} |\psi(x,t)|^2\, dx \tag{3.2}$$

and the Hamiltonian

$$\mathcal{H}[\psi] = \int_{-L}^{L} \Big(|\partial_x \psi|^2 - |\psi|^4 \Big)\, dx \tag{3.3}$$

are independent of t.

Proof. Identical to Theorem 2.1 of Chapter 2; see that proof for details. □

Remark 3.3. On a large but finite domain $(-L, L)$, the norm of the grey soliton is dominated by the background: $\mathcal{N} \approx 2A^2 L$. In our implementation we track the norm deviation $|\mathcal{N}_h(t) - \mathcal{N}_h(0)|$ rather than the absolute norm, since this quantity isolates the numerical error from the large constant background contribution.

The nonlinear Schrödinger and Gross–Pitaevskii equations govern some of the most precisely verified phenomena in physics: soliton propagation in optical fibres, matter-wave dynamics in Bose–Einstein condensates, modulational instability of nonlinear wavetrains, and the formation of rogue waves on the deep ocean. Both equations are completely integrable, and their solution manifold is rich — bright and dark solitons, breathers, cnoidal waves, and rational rogue-wave hierarchies.

This book constructs a rigorous computational atlas of those solutions. Fourteen chapters, each anchored to a closed-form exact solution, apply the Method of Lines: space is reduced to a high-order spectral operator, and the resulting differential system is integrated adaptively to pointwise accuracy of 10^{-10} or better. Conservation of the L^2 norm, momentum, and Hamiltonian energy is monitored as an independent diagnostic at every step. A closing chapter maps the limits of the pipeline and charts extensions toward non-integrable perturbations, stochastic forcing, and higher-dimensional collapse.

Dr. Þráinn Eiríksson is an applied mathematician who works where integrable systems meet high-accuracy scientific computing — on the numerical treatment of nonlinear wave equations in optics, cold-atom physics, and the study of rogue waves.

ISBN 13 978-831-05-3869-7
Paperback
9 788310 538697